



An Roinn Talmhaíochta,
Bia agus Mara
Department of Agriculture,
Food and the Marine

Published 23/11/2022

Avian Influenza Update

23rd November 2022

National Disease Control Centre

Number 17 of 2022

Highly Pathogenic Avian Influenza (HPAI H5N1) confirmed in a second poultry flock in Co. Monaghan

- On 22nd November, a second outbreak of highly pathogenic avian influenza (HPAI) subtype H5N1 was confirmed in a poultry flock (turkeys) in Co. Monaghan. This outbreak has occurred in an area very close the first outbreak (see Fig. 1).
- To mitigate the risk of spread of the disease, 3km protection and 10km surveillance zones have been introduced around the outbreak. This places additional controls on the movement of poultry and poultry products within the zones. Movements into and out of the zones are also subject to licensing. Additional surveillance measures also apply within the restricted zones. See figure 1 below for location of zones.
- An interactive map (for illustrative purposes) indicating the location of the protection and surveillance zones, courtesy of the Centre for Veterinary Epidemiology and Risk Analysis (CVERA), is available at the following link <https://arcg.is/14XyCO0>
- There remains a significant risk of further outbreaks and strict biosecurity is essential on all poultry and captive bird holdings
- All poultry and captive bird owners are reminded to **review and, where necessary, improve biosecurity procedures** on their premises to ensure that all appropriate measures are in place.




Fig. 1. Location of protection (red) and surveillance (green) zones around outbreak of HPAI in poultry in Co. Monaghan

- **Contact with wild birds and/or environmental contamination remain the major risk factors of HPAI incursion. Stringent biosecurity is the best defence!**
- **Flock owners should remain vigilant for any signs of disease in their flocks, and report any disease suspicion to their nearest Regional Veterinary Office**

Regulations in force: Housing order and mandatory biosecurity measures

- On the 7th November, new regulations (**Avian Influenza (Precautionary Confinement of Birds) Regulations 2022**) came into force which require flock keepers to confine all poultry and captive birds in their possession or under their control in a secure building. Where this is not possible, they must be secured in a manner that prevents access to other poultry, captive birds or wild birds. The Regulations may be found [here](#)
- **Housing birds is a support for biosecurity, not a replacement.** All flock owners are **legally required to maintain best practice biosecurity measures** to mitigate against the risk of introducing avian influenza into poultry and captive bird flocks since the introduction of Avian Influenza (Biosecurity measures) Regulations 2022 on 19th September 2022
- A copy of the [Avian Influenza \(Biosecurity measures\) Regulations 2022](#) and [Biosecurity measures Regulations accompanying guidance document](#) are also available on www.gov.ie/birdflu as well as further information on biosecurity

Risk of Highly Pathogenic Avian Influenza to poultry and captive birds

- Avian Influenza is **highly contagious for birds**. Due to the avian influenza surveillance carried out by DAFM, the disease is known to be circulating widely in wild bird populations, thereby presenting a risk of infection to Irish poultry and captive birds.
- The environment can become contaminated with virus through the faeces of infected birds. This presents a potential source of infection for poultry and captive birds
- **Biosecurity is the best defence** against the spread of disease, including avian influenza, to a poultry flock
- Members of the public are reminded **not to touch sick or dead wild birds**. Dogs should be kept on a lead when walking in areas where sick or dead wild birds are present
- **Poultry owners and those working with poultry** should beware of visiting areas, both at home and abroad, where deaths of wild birds have occurred (e.g.

coastal areas, bird colonies, etc.). Direct contact with sick or dead wild birds must be avoided

Avian influenza situation in wild birds

- On 18th November, HPAI H5N1 was confirmed in a peregrine falcon in Co. Offaly (see Fig. 2). This follows the confirmation of HPAI H5N1 in a mute swan in Co. Cavan on the 26th October and in pheasants in Co. Meath on the 11th November
- These cases of HPAI H5N1 demonstrates that the virus is present in wild birds in inland areas of Ireland, following cases found in wild birds in coastal areas since July
- An interactive map (for illustrative purposes) courtesy of the Centre for Veterinary Epidemiology and Risk Analysis (CVERA) displaying the location of avian influenza cases in wild and captive birds is available at the following link <https://arcg.is/19zHrf0>




Fig. 2. Locations of H5N1 avian influenza cases in wild birds and outbreaks in captive birds in Ireland since 1st July 2022

Avian Influenza situation in Northern Ireland

- The 3km protection and 10km surveillance zones implemented following the outbreak in a poultry flock in Co. Monaghan extend into Northern Ireland
- On 16th October, HPAI H5N1 was confirmed in captive birds at a Wildfowl and Wetlands Trust (WWT) near Strangford Lough. A 3km Captive Bird Monitoring

Zone has been implemented in the surrounding area to mitigate against onward disease spread

- This is the first outbreak of avian influenza in kept birds in Northern Ireland since last February
- On 17th October, an Avian Influenza Prevention Zone (AIPZ) was declared across Northern Ireland. The AIPZ places a legal requirement on all bird keepers in Northern Ireland to follow strict biosecurity measures.

Strict Biosecurity is Essential to Prevent Outbreaks

All poultry owners and keepers must implement very high levels of biosecurity on their holdings to help protect their flocks due to the current situation. All flock owners should ask themselves the following questions to help ensure you are implementing appropriate measures:

1. Who has access to the poultry areas?

Only allow **essential** persons access to the poultry houses e.g. husbandry, veterinary care. Ensure that dedicated clothing or disposable personal protective equipment (PPE) is worn including disposable gloves and overalls. Clean and disinfect boots before entering and immediately upon exiting each poultry house. Wash hands each time upon leaving a poultry house. Dedicated personnel should be appointed to manage each site

2. Have I checked the disinfectant points?

All disinfectant points must be maintained. **Read the label** on the disinfectant which is approved for use against avian influenza. Disinfectant footbaths must be changed regularly in accordance with the manufacturer's instructions. Contamination prevents the effective use of disinfectants. Footwear needs to be cleaned thoroughly before being disinfected in foot dips. **Wait the recommended contact time before entering the house.** Repeat checks with disinfectant points at entry and exit points to the premises

3. Can wild birds get access to the poultry?

Inspect the housing thoroughly. Ensure that there are no gaps where wild birds can enter the housing. Check the gutters and roof. Can contaminated rainwater enter the house? Can wild birds get access to or contaminate the drinking water system?

4. Is the feed and bedding stored securely?

Feed and bedding attract wild birds so secure storage of both is essential. If using straw, ensure it is either double wrapped and covered with a birdproof cover or stored in a shed with no wild bird access. Remove wrapping outside of the house.

Feed should be stored in a secure bin and any spills cleaned up immediately. Use wild bird deterrents

5. What do I do about feed delivery/egg collection vehicles that have to come onto the holding?

Disinfect wheels, wheel arches and underbody of all vehicles coming before allowing them to enter the premises and prior to leaving the premises. Do not allow any drivers to enter poultry houses. Ensure that drivers wear disposable gloves and disinfect footwear upon arrival (ensure that there is a disinfectant point near the feed bin/egg storage area)

6. What about equipment?

Any equipment that is used to transport poultry or poultry products must be thoroughly cleaned and disinfected prior to being used on the farm e.g., trolleys, trays, pallets, cages etc. Disposable single-use egg trays or plastic Keyes trays which are thoroughly cleaned and disinfected after each use can be used for egg collections. Don't forget about other equipment such as feeders and watering equipment and clean and disinfect equipment and housing between bird placements

7. Be vigilant

Check the feed and water intake and look for clinical signs of avian influenza. Report any suspicions immediately to DAFM and do not allow anyone on/off the premises pending investigation

Further information regarding Highly Pathogenic Avian Influenza H5N1

- The disease poses no food safety risk for consumers. Properly cooked poultry and poultry products, including eggs are safe to eat
- From October onwards each year we enter into the higher risk period for Avian Influenza in Ireland. Colder weather and decreasing daylight can extend the survival time of Avian influenza viruses in the environment
- This is due to the fact that migratory wild birds, the natural hosts of many avian influenza viruses, start to arrive in large numbers to overwinter here
- Migratory wild birds are known to carry the H5N1 subtype of Avian influenza, and this can increase the level of risk for a potential outbreak of Avian influenza in Ireland
- HPAI H5N1 is a notifiable disease both in Ireland and to the European Commission and the World Organisation for Animal Health (WOAH, formed known as OIE)

- Outbreaks of HPAI H5N1 in poultry in Ireland impact upon our international trade in poultry and poultry products
- The disease is highly contagious for birds, but risk of transmission to humans is considered very low.


DAFM surveillance for avian influenza

- Sick or dead wild birds can be reported to DAFM using the **Avian Check - Wild Bird App** <https://aviancheck.apps.rhos.agriculture.gov.ie>
- Alternatively, sick or dead wild birds may be reported to the local Regional Veterinary Office or contact the DAFM disease hotline on 01 607 2512 (during office hours) or 01 492 8026 (outside office hours)
- Where the Department has sufficient epidemiological surveillance information in particular geographical areas, it will not necessarily collect dead birds for testing
- DAFM has sufficient evidence of HPAI H5N1 in wild birds in coastal counties. As such, DAFM has stopped collecting birds for surveillance purposes from coastal counties. Surveillance continues as per current protocols in other counties (1 or more bird of prey-including owls, 3 or more waterfowl-including gulls, 5 or more of any other species of bird)
- Where dead wild birds are not required for surveillance purposes, and are present on public land, it is the responsibility of the local authority to safely dispose of the carcasses
- DAFM implements avian influenza surveillance measures which take place right throughout the year in both wild birds and poultry and form part of our well-established Early Warning System for the detection of avian influenza. The submission of dead wild birds which are on the target species list for avian influenza surveillance for testing, based on a risk assessment, at the Central Veterinary Research Laboratory forms part of this programme

Current Situation in poultry and captive birds in Europe

- HPAI continues to be reported in poultry and captive birds across the EU (see Fig. 3).
- Northern Ireland had six outbreaks of HPAI H5N1 in poultry during the 2021/2021 season. One outbreak of avian influenza in captive birds was reported on 11th October 2022

- Between 1st October 2022 and 20th November 2022 there have been , 117 outbreaks of HPAI H5N1 confirmed in England, 8 in Scotland and 3 in Wales
- The Animal and Plant Health Agency's (APHA) interactive map of restriction zones in England, Scotland and Wales is available to view [here](#)



No. of Highly Pathogenic Avian Influenza reports to ADIS per week

Fig 3. Number of confirmed outbreaks and cases of HPAI in poultry, captive birds and wild birds in EU Member States during the period 19th September-22nd November 2022 (Source: [Istituto Zooprofilattico Sperimentale delle Venezie](#))

Current wild bird situation in the rest of Europe

- HPAI H5N1 is currently circulating in wild birds, especially breeding seabirds around the UK's coast
- Large die-offs of several species of colonial breeding birds associated with avian influenza have been reported recently in the Netherlands, Scotland, Iceland, Norway, France, and Greece ([EFSA, 2022](#))
- DAERA have produced an interactive map viewer displaying locations of cases of avian influenza in wild birds in Northern Ireland which is available to view [here](#)
- Over 95% of cases in wild birds have been confirmed to be of subtype H5N1, with small numbers of H5N3 and H5N8 also confirmed




Fig. 4 HPAI detections in wild and captive during the period 01 October 2022 – 18 November 2022

What can poultry owners do to reduce the risk of avian influenza?

Flock owners should **remain vigilant for any unexplained signs of disease** in their flocks, and report any disease suspicion to their nearest [Regional Veterinary Office](#)

1. Be vigilant- Monitor your birds carefully for any clinical signs of disease. If in doubt, contact your veterinary practitioner for advice. Signs of avian influenza include:

- Death/high mortalities in a flock
- Depression/lethargy
- Loss of appetite
- Respiratory distress such as gaping beak, coughing, sneezing, gurgling, rattling
- Swelling and blue discolouration of combs, wattles, neck and throat
- Diarrhoea
- Reduced egg production or no egg production

2. Biosecurity

A copy of the Avian Influenza (Biosecurity measures) Regulations 2022, accompanying guidance document and additional species specific biosecurity leaflets can be found [here](#).

Personnel and Visitors

- Wear disposable or dedicated outer clothing when in contact with your birds
- Wear boots that can be disinfected, disposable gloves and wash hands after contact with poultry
- Only allow essential personnel access to your birds

Buildings and Equipment

- Regularly clean and disinfect hard surfaces such as paths and walkways
- Provide clean boot washing and disinfection facilities at all entrances/exits and ensure that people use them
- Clean and disinfect any vehicles or equipment that comes into contact with the birds after use
- Have suitable pest control in place

Wild Birds

- Feed and water birds inside or at least under cover
- Clean up any waste feed in outdoor areas frequently
- Use netting or wild bird decoys or deterrents to keep wild birds away from poultry areas

Management

- Keep ducks and geese separate to other poultry species
- The Department also reminds all poultry owners, including those who keep only 1 or 2 birds of their legal obligation to register their premises with the Department.
- Use an [approved disinfectant](#) against Avian influenza in accordance with the manufacturer's instructions

Further information:

DAFM has a wide variety of biosecurity resources for avian influenza- please see:

<https://www.gov.ie/en/publication/7e230-poultry/>

To register a poultry flock:

<https://www.gov.ie/en/service/984df6-poultry-register/>

RVO contact details:

[Regional Veterinary Office](#)

List of target species of wild birds for avian influenza

surveillance: [List of Target Species for Avian Influenza](#)

[Surveillance](#)

European Commission

https://ec.europa.eu/food/animals/animal-diseases/diseases-and-controlmeasures/avianinfluenza_en

World Organisation for Animal Health:

[Avian Influenza - WOA - World Organisation for Animal Health](#)

European Food Safety Authority:

<https://www.efsa.europa.eu/en/topics/topic/avian-influenza>

National Disease Control Centre

23rd November 2022